

Math 111 Problem Set 2 Solutions

Edward Doolittle

October 20, 2005

1. (a) $2\log_{10} 3 - \log_{10} 90 = \log_{10} 3^2 - \log_{10} 90 = \log_{10} 9 - \log_{10} 90 = \log_{10}(9/90) = \log_{10} 10^{-1} = -1$.
(b) $\ln(e^{2^3}) = 2^3 \ln(e) = 8 \cdot 1 = 8$.

2. (a) Exponentiating both sides,

$$\begin{aligned}e^{\ln(5x-2)} &= e^{-7} \\5x - 2 &= e^{-7} \\5x &= e^{-7} + 2 \\x &= \frac{1}{5}(e^{-7} + 2).\end{aligned}$$

My calculator tells me that $x = 0.4002$ to 4 decimal places.

- (b) Taking the log base 2 of both sides,

$$\begin{aligned}\log_2(2^{3x+4}) &= \log_2(10) \\(3x + 4)\log_2(2) &= \log_2(10) \\3x + 4 &= \log_2(10) \\3x &= \log_2(10) - 4 \\x &= \frac{1}{3}(\log_2(10) - 4)\end{aligned}$$

Since there is no \log_2 button on my calculator I have to use the chain rule to evaluate $\log_2(10) = \ln(10)/\ln(2)$; with that, my calculator tells me $x = -0.2260$.

3. (a) Write $f(x) = (x \ln x)^{1/2}$. Then by the chain rule and product rule,

$$\begin{aligned}f'(x) &= \frac{1}{2}(x \ln x)^{-1/2} \frac{d}{dx}(x \ln x) \\&= \frac{1}{2}(x \ln x)^{-1/2} \left(\left(\frac{d}{dx} x \right) \ln x + x \frac{d}{dx} \ln x \right) \\&= \frac{1}{2}(x \ln x)^{-1/2} \left(\ln x + x \frac{1}{x} \right) \\&= \frac{1}{2}(x \ln x)^{-1/2} (\ln x + 1).\end{aligned}$$

- (b) Rewrite 2^u in terms of the (natural) exponential function: $2^u = (e^{\ln 2})^u = e^{(\ln 2)u}$. Then by the chain rule,

$$\begin{aligned}g'(\theta) &= \frac{d}{d\theta} e^{(\ln 2) \sin \theta} \\&= e^{(\ln 2) \sin \theta} \frac{d}{d\theta} ((\ln 2) \sin \theta) \\&= 2^{\sin \theta} (\ln 2) \cos \theta.\end{aligned}$$

4. (a) Take the logarithm of both sides:

$$\begin{aligned}\ln y &= \ln \left((2x^3 + 1)^4 (x - 1)^{-1/2} \right) \\ &= 4 \ln(2x^3 + 1) - \frac{1}{2} \ln(x - 1)\end{aligned}$$

Then by implicit differentiation on the left side and the chain rule on the right side,

$$\begin{aligned}\frac{y'}{y} &= 4 \frac{1}{2x^3 + 1} \frac{d}{dx}(2x^3 + 1) - \frac{1}{2} \frac{1}{x - 1} \frac{d}{dx}(x - 1) \\ &= \frac{24x^2}{2x^3 + 1} - \frac{1}{2x - 2}\end{aligned}$$

Multiplying both sides by y and substituting what we know about y ,

$$\begin{aligned}y' &= y \left(\frac{24x^2}{2x^3 + 1} - \frac{1}{2x - 2} \right) \\ &= (2x^3 + 1)^4 (x - 1)^{-1/2} \left(\frac{24x^2}{2x^3 + 1} - \frac{1}{2x - 2} \right).\end{aligned}$$

Further simplification is possible, but not necessary unless the question asks for it.

- (b) Take the logarithm of both sides:

$$\begin{aligned}\ln h &= \ln \left(\frac{\sin^2 x \cos^4 x}{(x^2 + 1)^3} \right) \\ &= 2 \ln \sin x + 4 \ln \cos x - 3 \ln(x^2 + 1).\end{aligned}$$

Differentiate the above equation (by implicit differentiation on the left side and the chain rule on the right side):

$$\begin{aligned}\frac{h'}{h} &= 2 \frac{1}{\sin x} \cos x + 4 \frac{1}{\cos x} (-\sin x) - 3 \frac{1}{x^2 + 1} (2x) \\ &= \frac{2 \cos x}{\sin x} - \frac{4 \sin x}{\cos x} - \frac{6x}{x^2 + 1}.\end{aligned}$$

Multiplying by h ,

$$h'(x) = \frac{\sin^2 x \cos^4 x}{(x^2 + 1)^3} \left(\frac{2 \cos x}{\sin x} - \frac{4 \sin x}{\cos x} - \frac{6x}{x^2 + 1} \right).$$

5. Differentiating both sides of the equation, then applying the chain rule,

$$\begin{aligned}\frac{d}{dx} y^2 &= \frac{d}{dx} e^{x^2 + y^2} \\ 2y \frac{d}{dx} y &= e^{x^2 + y^2} \frac{d}{dx} (x^2 + y^2) \\ 2yy' &= e^{x^2 + y^2} (2x + 2yy').\end{aligned}$$

Solving for y' ,

$$\begin{aligned}2yy' - 2ye^{x^2 + y^2} y' &= 2xe^{x^2 + y^2} \\ 2y(1 - e^{x^2 + y^2})y' &= 2xe^{x^2 + y^2} \\ y' &= \frac{2xe^{x^2 + y^2}}{2y(1 - e^{x^2 + y^2})} \\ &= \frac{xe^{x^2 + y^2}}{y(1 - e^{x^2 + y^2})}.\end{aligned}$$

6. Let I be the required result. First expand the integrand using the formula $(a + b)^2 = a^2 + 2ab + b^2$:

$$I = \int_4^{16} \left(\sqrt{x} + \frac{3}{\sqrt{x}} \right)^2 dx = \int_4^{16} \left(x + 6 + \frac{9}{x} \right) dx.$$

Integrating each term,

$$\begin{aligned} I &= \int_4^{16} x dx + \int_4^{16} 6 dx + \int_4^{16} \frac{9}{x} dx \\ &= \frac{x^2}{2} \Big|_4^{16} + 6x \Big|_4^{16} + 9 \ln x \Big|_4^{16} \\ &= 128 - 8 + 96 - 24 + 9 \ln(16) - 9 \ln(4) \\ &= 192 + 18 \ln(2). \end{aligned}$$

To four decimal places, $I = 204.4766$.

7. (a) To find the inverse, solve for the independent variable t :

$$\begin{aligned} Q &= Q_0(1 - e^{-t/a}) \\ \frac{Q}{Q_0} &= 1 - e^{-t/a} \\ 1 - \frac{Q}{Q_0} &= e^{-t/a} \\ \ln(1 - Q/Q_0) &= \ln(e^{-t/a}) = -t/a \\ -a \ln(1 - Q/Q_0) &= t, \end{aligned}$$

so the inverse function is

$$Q^{-1}(q) = -a \ln(1 - q/Q_0).$$

$Q^{-1}(q)$ should be interpreted as the amount of time it takes for the camera to reach charge level q .

- (b) By the previous result, we need to find $Q^{-1}(0.75Q_0)$:

$$\begin{aligned} Q^{-1}(0.75Q_0) &= -3 \ln(1 - (0.75Q_0)/Q_0) \\ &= -3 \ln(1 - 0.75) \\ &= -3 \ln(1/4) \\ &= 6 \ln(2). \end{aligned}$$

Calculator says $t = 4.2$ seconds approximately.

8. The slope of the line $x + 2y = 5$ is $-1/2$. The slope of any perpendicular line is the negative reciprocal, i.e., 2. We need to find a line tangent to $y = e^x$ with slope 2. The line tangent to the curve at the point (x, e^x) has slope $\frac{d}{dx} e^x = e^x$, so we need to solve the equation $e^x = 2$: $x = \ln(2)$. At the point with the given x -value has coordinates $(\ln(2), e^{\ln(2)}) = (\ln(2), 2)$. The point-slope form of the tangent line is $y - 2 = 2(x - \ln(2))$.
9. Using the hint and the rules for exponents,

$$\begin{aligned} \lim_{x \rightarrow 0^+} x^{\ln x} &= \lim_{x \rightarrow 0^+} (e^{\ln x})^{\ln x} \\ &= \lim_{x \rightarrow 0^+} e^{(\ln x)^2} \\ &= e^{\lim_{x \rightarrow 0^+} (\ln x)^2}. \end{aligned}$$

As $x \rightarrow 0^+$, $\ln x \rightarrow -\infty$, so $(\ln x)^2 \rightarrow \infty$. Therefore $\lim_{x \rightarrow 0^+} x^{\ln x} = e^\infty = \infty$.

10. Here are two different solutions:

(a) By formula 8 of section 7.4 and the continuity of the power function $f(x) = x^a$,

$$\lim_{n \rightarrow \infty} \left(1 + \frac{a}{n}\right)^n = \lim_{n \rightarrow \infty} \left(\left(1 + \frac{a}{n}\right)^{n/a}\right)^a = \lim_{h \rightarrow 0^+} \left((1+h)^{1/h}\right)^a = \left(\lim_{h \rightarrow 0^+} (1+h)^{1/h}\right)^a = e^a$$

where $h = a/n$.

(b) By the continuity of the natural logarithm function,

$$\ln \lim_{n \rightarrow \infty} \left(1 + \frac{a}{n}\right)^n = \lim_{n \rightarrow \infty} \ln \left(1 + \frac{a}{n}\right)^n = \lim_{n \rightarrow \infty} n \ln(1 + a/n) = \lim_{n \rightarrow \infty} \frac{\ln(1 + a/n)}{1/n} = a \lim_{n \rightarrow \infty} \frac{\ln(1 + a/n)}{a/n}.$$

Letting $h = a/n$ the limit becomes

$$a \lim_{n \rightarrow \infty} \frac{\ln(1 + a/n)}{a/n} = a \lim_{h \rightarrow 0^+} \frac{\ln(1 + h)}{h} = a \cdot 1 = a$$

by the result of problem 87 of section 7.4. Therefore

$$\ln \lim_{n \rightarrow \infty} \left(1 + \frac{a}{n}\right)^n = a$$

and exponentiating both sides,

$$\lim_{n \rightarrow \infty} \left(1 + \frac{a}{n}\right)^n = e^a$$

as required.