

MATH122 200610 Quiz 3B Solutions

Edward Doolittle

Wednesday, March 1, 2006

1. We must find a matrix B such that

$$\begin{bmatrix} 2 & -3 \\ -4 & 6 \end{bmatrix} \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

Splitting B and O into columns, we have two conditions

$$\begin{bmatrix} 2 & -3 \\ -4 & 6 \end{bmatrix} \begin{bmatrix} b_{11} \\ b_{21} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

and

$$\begin{bmatrix} 2 & -3 \\ -4 & 6 \end{bmatrix} \begin{bmatrix} b_{12} \\ b_{22} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

The two conditions are the same, except for the subscripts, so we concentrate on solving the first. The condition is a linear system which can be solved by row reduction. The system corresponds to the augmented matrix

$$\left[\begin{array}{cc|c} 2 & -3 & 0 \\ -4 & 6 & 0 \end{array} \right].$$

Adding 2 times row 1 to row 2 gives

$$\left[\begin{array}{cc|c} 2 & -3 & 0 \\ 0 & 0 & 0 \end{array} \right].$$

The system is in row echelon form. Multiplying the first row through by $1/2$ gives

$$\left[\begin{array}{cc|c} 1 & -3/2 & 0 \\ 0 & 0 & 0 \end{array} \right].$$

The system has solution $b_{11} = 3s$, $b_{21} = 2s$. Taking $s = 1$, for example, we obtain the solution $b_{11} = 3$, $b_{21} = 2$. (We can't take $s = 0$ because B can't have any zero entries, but any other value for s would work.) The solution for the second column of B is exactly the same; for variety we can take $s = -1$, for example, and obtain $b_{12} = -3$, $b_{22} = -2$. This gives

$$B = \begin{bmatrix} 3 & -3 \\ 2 & -2 \end{bmatrix}$$

as one example of a B that works. Checking by matrix multiplication,

$$\begin{aligned} AB &= \begin{bmatrix} 2 & -3 \\ -4 & 6 \end{bmatrix} \begin{bmatrix} 3 & -3 \\ 2 & -2 \end{bmatrix} \\ &= \begin{bmatrix} 2(3) - 3(2) & 2(-3) - 3(-2) \\ -4(3) + 6(2) & -4(-3) + 6(-2) \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}. \end{aligned}$$

2. To find x_1 , x_2 , x_3 we must solve the system $B\mathbf{x} = \mathbf{0}$. Forming the augmented matrix,

$$\left[\begin{array}{ccc|c} -1 & 2 & 8 & 0 \\ 1 & 0 & -2 & 0 \\ 2 & 4 & 8 & 0 \end{array} \right].$$

Adding 1 times row 1 to row 2 and 2 times row 1 to row 3 we obtain

$$\left[\begin{array}{ccc|c} -1 & 2 & 8 & 0 \\ 0 & 2 & 6 & 0 \\ 0 & 8 & 24 & 0 \end{array} \right].$$

Adding -4 times row 2 to row 3 we obtain

$$\left[\begin{array}{ccc|c} -1 & 2 & 8 & 0 \\ 0 & 2 & 6 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right].$$

The system is now in row echelon form. Multiplying row 1 by -1 and row 2 by $1/2$ we obtain

$$\left[\begin{array}{ccc|c} 1 & -2 & -8 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right].$$

Adding 2 times row 2 to row 1 we finally obtain

$$\left[\begin{array}{ccc|c} 1 & 0 & -2 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right].$$

The system is in reduced row echelon form, and we can read the solution directly from it:

$x_1 = 2s$, $x_2 = -3s$, $x_3 = s$. Taking $s = 1$, for example, we get the non-trivial linear relation $2\mathbf{b}_1 - 3\mathbf{b}_2 + \mathbf{b}_3 = \mathbf{0}$. (You should double check that that linear relation applies to the \mathbf{b}_i s.)

Now, let $C = AB$. By matrix multiplication,

$$C = \begin{bmatrix} 2 & 3 & 1 \\ -2 & -3 & 0 \\ 2 & 1 & -1 \end{bmatrix} \begin{bmatrix} -1 & 2 & 8 \\ 1 & 0 & -2 \\ 2 & 4 & 8 \end{bmatrix} \\ = \begin{bmatrix} 3 & 8 & 18 \\ -1 & -4 & -10 \\ -3 & 0 & 6 \end{bmatrix}.$$

It is straightforward to check that the same relationship as held for the \mathbf{b}_i s also holds for

the \mathbf{c}_i s, i.e., $2\mathbf{c}_1 - 3\mathbf{c}_2 + \mathbf{c}_3 = \mathbf{0}$:

$$2\mathbf{c}_1 - 3\mathbf{c}_2 + \mathbf{c}_3 \\ = 2 \begin{bmatrix} 3 \\ -1 \\ -3 \end{bmatrix} - 3 \begin{bmatrix} 8 \\ -4 \\ 0 \end{bmatrix} + \begin{bmatrix} 18 \\ -10 \\ 6 \end{bmatrix} \\ = \begin{bmatrix} 6 \\ -2 \\ -6 \end{bmatrix} + \begin{bmatrix} -24 \\ 12 \\ 0 \end{bmatrix} + \begin{bmatrix} 18 \\ -10 \\ 6 \end{bmatrix} \\ = \begin{bmatrix} 6 - 24 + 18 \\ -2 + 12 - 10 \\ -6 + 0 + 6 \end{bmatrix} = \mathbf{0}.$$

(As a short cut, you know the relationship must hold because $(AB)\mathbf{x} = A(B\mathbf{x}) = A\mathbf{0} = \mathbf{0}$ by the associativity of matrix multiplication.)